

PERCEPTION OF UNIVERSITY STUDENTS IN PALESTINE REGARDING PREVENTIVE MEASURES DURING COVID-19 PANDEMIC

MUSTAFA MOHAMMAD SHOULI

Assistant Professor, Nablus University for Vocational & Technical Education.
Corresponding Author Email: Mustafa.shouli@nu-vte.edu.ps

KHAILA MOHAMMAD SHOULI

Lecturer, Nablus University for Vocational & Technical Education. Email: Khaila.shouli@nu-vte.edu.ps

BILAL MOHAMMAD ISSA

Lecturer, Nablus University for Vocational & Technical Education, Email: bilal.issa@nu-vte.edu.ps

Abstract

Knowledge, perception, and preventive measures with regard to COVID-19 among the university student are of immense importance and are critical for the community prevention/ control of this pandemic. Aim to assess the perception of university students in Palestine regarding preventive measures during COVID-19 pandemic. Research design: A cross-sectional survey consisted of 2094 university students. Setting: the study was conducted at Modern University College, An-Najah National University, Palestine Polytechnic University, Al-Quds Open University, Ibn Sina College of Health Sciences and Arab American of University /West Bank / Palestine. The data collected was from the participant university students and started in first October 2020 to end of December 2020. Subjects: A convenience sample with different Bachelor degree programs. Tools: A self-designed questionnaire was used for data collection and included the following parts. 1: demographic particulars of the university students, II. University students Knowledge are regarding Coronavirus, isolation for patient with Covid-19 at home, also their knowledge regarding following for preventive measures outside home from [15; [16], III: preventive measure as practices were assessed for university students outside home Part IV: assessed University student perception toward Covid-19. Results: the current study reveals that 80.2% of the studied sample of students had a satisfactory total knowledge regarding preventive measures outside home, 91% of them had a satisfactory total knowledge regarding preventive measures inside home among patient with Covid-19, and 95.5% of them had a satisfactory total knowledge regarding Covid-19 disease. There was a statistically highly significant relation between total knowledge and total practices outside home (p -value<0.001), and statistically significant relation between total knowledge and total perception (p -value0.005).Conclusion: based on the current study, students were observed to have substantial knowledge, practices for preventive measures, and a positive perception toward COVID-19 pandemic. Recommendation: It is the responsibility of the state to take all preventive measures for the safety of the community and public. There is an urgent need for research to address burden of the Covid-19 pandemic on college students.

INDEXTERMS: Perception, University Students, Preventive Measures, COVID-19 Pandemic,

INTRODUCTION:

Coronavirus disease 2019 which is abbreviated "COVID-19" is a developing respiratory disease that is triggered by a new coronavirus. The newest member of the coronavirus family (COVID-19) has been recently identified as acute and severe respiratory syndrome in humans. It was established that in China, 18.5% of the patients with COVID-19 developed to the severe stage, which is characterized by acute respiratory distress syndrome, septic shock, difficult-to-tackle metabolic acidosis, and bleeding and

coagulation dysfunction [24]. Some of the unparalleled measures adopted to control the COVID-19 transmission in Hubei and other provinces of China and other parts of the world, including the suspension of public transportation, the closing of public places, management of communities, and isolation and care for infected people and suspected cases [23].

The symptoms of COVID-19, some infected people may not experience any symptoms, whereas others may have mild symptoms. Around 80% of patients recover without needing any special treatment, while one out of six patients develop more severe symptoms, not limited to difficulty breathing. These patients are most likely to be elderly or with underlying medical conditions, such as high blood pressure, heart problems or diabetes [23].; [21]. Additionally, young people are not invincible against coronavirus; world data have shown that a large number of the patients were young, among which many needed hospitalization [5] ; [6]; [8].

The pandemic coronavirus disease (COVID-19) is a highly infectious disease originated from Wuhan city of China and is still swiftly spreading and infecting public all over the world. It is caused by a virus known as severe acute respiratory syndrome coronavirus (SARS-CoV-2) [18] .On December 31st of 2020, WHO declared this outbreak as a Public Health Emergency of International Concern and publicized a name for the new coronavirus disease as Covid-19 and on 11th of March 2020, WHO confirmed Covid-19 as pandemic.

Globally, a total of 8,236,326 cases were identified with 40,598 new cases in more than 170 countries till 1st April 2020. The European region is more affected then the rest of the continents. About 464,212 cases were reported with 30,089 deaths in European region, followed by Regions of America with 188,751 cases along with 3,400 causalities. The cases of Western Pacific region, Eastern Mediterranean region, South-East Asia region and African regions are 106,422 cases (3,701 deaths), 54,281 cases (3,115 deaths), 5,175 (195 deaths) and 4,073 cases (91 deaths) respectively. Highest no. of cases was reported in United State (163,199 cases), followed by Italy, Spain, China, France and Germany. However, the mortality rate is greater in Italy (11.7%), followed by Spain (8.7%), United Kingdom (7.11) and France (6.8%). As a result of traveling, an outbreak occurred in several countries especially the European regions are more affected than the rest of world [25] .

Perception is essential to understanding their response, behaviour, and adoption of individual-level preventive measures (e.g. wearing masks, washing hands) in case of an infectious disease outbreak and its aftermath. Identifying risk perception will not only help mitigate the devastating mortality and morbidity burden, but also economic loss. With the relatively long period needed for the majority of the population to be vaccinated against COVID-19 and the emergence of new COVID-19 variants , understanding and addressing behaviour to reduce transmission and spread of infection is imperative [31].To avoid further spikes of new cases and unavoidable enforcement of lockdowns . Such insight will help identify population groups with relatively low risk perceptions or low adoption of

preventive measures and enable the design of policies and interventions tailored to these populations [32].

The use of face masks and hand sanitizers, a ban on social events, and working from home, etc., were among the measures that were followed to curb the spread of COVID-19. Taiwanese people panic-bought nearly all available mask and hand sanitizer stocks within 2 weeks of the first case in Taiwan (21 January 2020). Movie theatres, restaurants, and malls had diminishing crowds [22].

The perception and knowledge of health literacy skills permit students to manage their own well-being by improving their communication with doctors, and making smart healthcare decisions. Nevertheless, public adherence to control Covid-19 is influenced greatly by their knowledge, perception, and their adherence for preventive measures. Therefore, the present study was aimed to explore the perception of university students in Palestine regarding preventive measures during COVID-19 pandemic [20].

Significance of the study:

According to the Palestinian Ministry of Health (MOH), the first cases had been detected at a hotel in Bethlehem area, where a group of Greek tourists had visited the hotel in late February 2020, with two later diagnosed with the virus [28].According to The United Nations Relief and Works Agency (UNRWA), the prolonged blockade imposed has severely impacted the socioeconomic and health conditions of the Occupied Palestinian Territories and Gaza strip [30].The longstanding movement restrictions have undermined Gaza's economy and the checkpoints in West Bank to high levels of unemployment, food insecurity, aid dependency, and poor standards of hygiene and sanitation [29]. Consequently, the government response was swift and immediate, started by launching a social media campaign encouraging people to stay at home and to follow the ministry of health instructions.

METHODS:

i. Technical design:

Aim: this study aimed to explore the perception of university students in Palestine regarding preventive measures during COVID-19 pandemic.

Research Design: A cross-sectional sample survey consisted of 2094 university students for male and female ,aged from 18 >20 years by Using online electronic data collection, the survey method was used because the study aimed to assess the knowledge level, perception , practices and their preventive measures of COVID -19 among Palestinian University Students.

Setting: The study was conducted at six universities in West Bank / Palestine was Modern University College, An-Najah National University, Palestine Polytechnic University, Al-Quds Open University, Ibn Sina College of Health Sciences and Arab American of University.

Sample: A convenience sample included 2094 university students with different Bachelor degree programs, from six previous mentioned universities and divided into 965 from health sciences, 382 financial and administrative sciences, 249 engineering and 498 participated from social sciences. All participants responded to the online Google form link sent to what's-App groups. The study was conducted between November and December, 2020. The University students used a Google form shared through email and mobile messages.

Tools for data collection:

Tool I.: A self-designed questionnaire was used for data collection and included the following parts:

A.-: Demographic particulars of the participant. The demographic variables consist of age, gender, and specialty, place of residence, family income and number of family.

B. Knowledge was assessed for university student using a 20-item questionnaire adapted from [15; [16]. slightly customized to suit college students. The items were attributed to knowledge about Coronavirus pandemic disease (10 Questions), knowledge among isolation for patient with Covid-19 at home (7 Questions), knowledge regarding following for preventive measures outside home (3 Questions). 3-point scale were fixed as complete correct= (3), incomplete correct = (2) and incorrect= (1). The total score was recorded as satisfactory > 85%, unsatisfactory < 85%. All knowledge questions were based on Centers for Disease Control and Prevention (CDC) fact sheets.

Part III: preventive measures were assessed for practices university students outside home like hand washing, mask wearing, avoiding crowded places; keeping social distance and avoiding unnecessary travelling etc. The questionnaire was constructed on the basis of the published literature from the [25].

(<https://main.icmr.nic.in/>) for prevention of SARS-Cov2 transmission. The responses were fixed as always= (2), occasional= (1), and never= (0) respectively. The total score was recorded as satisfactory > 85%, unsatisfactory < 85%.

Part IV: Assessed the University students' perception toward Covid-19: approximately eleven items from a previous survey were used to assess the student's perception. The questions were related to agree that successfully controlled in elimination of Covid-19 changed for someone if affected with Covid-19. Do you: ask for consult from responsible authorities in the event of emergency symptoms, think the local authorities will succeed in reducing the spread of disease, think the preventive measures of the government are early, think the preventive measures of the government were enough, think you do not get off the Covid-19, think it is not necessary to take precautionary measures for children and older. Isolation of affected people is an effective way to reduce the spread of the Covid-19, and early treatment is the effective way to control the prevalence of Covid-19 as well. The scoring choice is the option "agree" which added two points, the option "not agree" added one, while zero was added for the "don't know" option in each behaviour-related question.

II- Operational Design:

The operational design was included; preparatory phase, ethical considerations, tools validity and reliability, pilot study and field of work.

Preparatory phase:

The researcher reviewed the local and international literature to be aware of various aspects of the research problem, by using books, journals and internet search and to design the study tools.

Content Validity & reliability:

Content validity was done to identify the degree to which the tools measure what was supposed to be measured. The translated tools were examined by a panel of three experts in the field of community who agreed that it is valid and relevant with the aim of the study. Internal consistency was measured to identify the extent to which the items of the tools measure the same concept and the extent to which the items are correlated with each other. The internal consistency which estimated reliability by Cronbach's Alpha was 0.862.

Pilot study: A pilot study was carried out on 10% (209) university student from the total number of sample to assess the tools' clarity, objectivity and feasibility, and to estimate the time needed for data collection. Those students in the pilot study were included in the main study sample since some modifications were made.

Statistical analysis: descriptive statistical measures including frequency, percentage, and mean score were used to report the findings. The chi-square test was used to determine the association between the categories. Statistical analysis was performed using SPSS (IBM SPSS Statistics for Windows, Version 25.0; IBM Corp., Armonk, NY). P-value < 0.05 was considered not statistically significant & P-value < 0.001 was considered statistically highly significant.

Ethical consideration: Ethical approval was obtained from the previous six mentioned Universities to conduct the study. All participants provided informed consent before participating in the study by using a Google form shared through email, mobile messages, and in print with students from the previous mentioned setting.

III. RESULT:

Table 1: Distribution of university students in Palestine according to their demographic characteristics (n= 2094)

Demographic Characteristics	No	%
Age(years)		
• 18- <19	420	20.1
• 19- <20	670	32.0
• >20	1004	47.9
Mean± SD= (19.56±3.069)		
Gender		
• Male	1119	53.4
• Female	975	46.6
Place of residence		
• City	564	26.9
• Village	967	46.2
• Camps	563	26.9
Family income		
• Enough	1920	91.7
• Not enough	174	8.3
Number of family		
• 2 - 4	743	35.5
• 5 – 7	1000	47.8
• > 7	351	19.7
University name		
Modern University College	606	28.9
Arab American University	331	15.8
An-Najah National University	297	14.2
Palestinian Polytechnic University	308	14.7
Ibn-Sina College for Nursing and Midwife	224	10.8
Al-Quds Open University	328	15.6
Programs		
Health Sciences	965	46.1
Financial and Administrational Sciences	382	18.2
Engineering	249	11.9
Social Sciences	498	23.8

Table 1: A total of 2094 students completed the online questionnaire. The mean/average age was (20.56±3.069) years and 53.3% were male. 46.1 Health Sciences, 18.2 financial and administrative Sciences, 11.9 engineering, 23.8 Social Sciences. Regarding residence, results show that 46.2% resided in village, while 26.9% in city and campus

residents. 47.8% of the students live with their family consisting of 5-7members. Additionally, 91.7% of the students have enough family income.

Table 2: Distribution of university students in Palestine according to their knowledge regarding Covid-19 pandemic (n=2094)

Items	Correct		Incomplete correct		Incorrect	
	N	%	N	%	N	%
*Meaning of Covid-19	2074	99.0	20	1.0	0	0.0
*Causes of Covid-19	2016	96.3	78	3.7	0	0.0
*Methods of transmission	2094	100.0	0.0	0.0	0.0	0.0
*Signs & symptoms of disease	1858	88.7	224	10.7	12	0.6
*How can you protect yourself and others from infection	1787	85.3	202	9.6	105	5.1
*Time for repetition tests of abnormal results	1179	56.3	242	11.6	673	32.1
*Medical tests to discover the virus of Covid-19	1526	72.8	482	27.2	86	4.0
*Prophylactic anticoagulation drugs used	1542	73.6	368	17.6	184	8.8
* Time to give anticoagulant to prevent blood Clotting	1359	64.9	490	23.4	245	11.7
*Treatment of Covid-19	1963	93.7	131	6.3	0	0.0

Table 2: shows that the great majority of the studied students (99%, 96% & 93.7) have a correct answer about the meaning of covid-19, its causes & the treatment respectively. All of them have a correct answer about transmission of Covid; also, the majority (88.7& 85.3) have correct answers regarding signs and symptoms & protection methods from infection respectively. While, 56.3% of them know the time for repetition tests, 11.6% have incomplete answers, and 32.1% did not know the time. Regarding test to discover the virus & anticoagulation drugs 72.8% & 73.6% have correct answers, 27.2% &17.6% have incomplete answers, and only 4%& 8.8% did not know, respectively. In addition, 64.9 have correct answers, 23.4% have incomplete answers & 11.7% have incorrect answers about timing to give anticoagulation.

Table 3: Distribution of university students in Palestine according to their knowledge among isolation for patient with Covid-19 at home (n=2094)

Items	Correct		Incomplete correct		Incorrect	
	N	%	N	%	N	%
*What is patient isolation at home	1994	95.2	100	4.8	0	0.0
*Meaning of self-quarantine	2071	98.9	23	1.1	0	0.0
*Type of medication during home isolation	1957	93.5	115	5.5	22	1.0
* Information about Precautions measures for patients with Covid-19 isolation	1981	94.6	100	4.8	13	0.6
* Information to deal with visitors for patients with Covid-19 isolation	1954	93.3	100	4.8	40	1.9
*Information for environmental cleaning to prevent Covid-19	1883	89.9	177	8.4	34	1.6
* Who are the preventive measures to disinfect and clean upholstery to prevent infection with covid-19?	1647	78.7	315	15.0	132	6,3

Table 3: reveals that the great majority (95.2%, 98.9%, 93.5 & 94.6%) respectively of studied students have correct answers regarding the determination of the patient who needs isolation at home, meaning self-quarantine, type of medication used during isolation, precaution with visitors, and Precautions measures for patients with Covid-19 isolation. Furthermore, 89.9% & 78.7% respectively have correct answers regarding information for environmental cleaning to prevent Covid-19 & the precautionary measures to disinfect and clean upholstery to prevent infection with covid-19. While, limited number of them have incomplete & incorrect answers towards precautionary measures to deal with a patient with Covid-19 at home.

Table 4: Distribution of university students' knowledge regarding followed preventive Measures outside home (n=2094)

Items	Correct		Incomplete correct		Incorrect	
	N	%	N	%	N	%
* The preventive measures that must be followed to avoid close contact with others	1908	91.1	124	5.9	62	3.0
* The preventive measures for respiratory hygiene to protect from Covid-19	1769	84.5	232	11.0	93	4.5
* What is personal protective equipment used to protect from covid-19	1750	83.6	243	11.6	101	4.8

Table 4: clarifies that 91.1% of studied students have correct answers regarding followed the precautionary measures outside home and 4.5% have incorrect answers regarding followed the precautionary measures for respiratory hygiene to protect from Covid-19. Also, 83.6% of them have correct answers regarding using personal protective equipment.

Figure 1: Distribution of studied university students according to their total satisfactory knowledge (n=2094)




Figure 1: illustrates that 80.2% of the studied students have a satisfactory total knowledge regarding followed the preventive measures outside home, 91% of them had a satisfactory total knowledge regarding isolation of patient affected with Covid-19 inside home while 95.5% of them have a satisfactory total knowledge regarding Covid-19 disease.

Table 5: Distribution of university students according to their practices for the preventive Measures regarding Covid-19 disease outside home (n=2094)

Items	Always		Occasional		Never	
	N	%	N	%	N	%
*Avoiding being in crowded places	1788	85.4	230	11.0	76	3.6
*Wearing the face mask when leaving the house	1240	59.2	820	39.2	34	1.6
*Practicing carefully cleaned hands using an antiseptic	1197	57.2	710	33.9	187	8.9
*Using tissue papers to cover mouth and nose during sneezing or coughing	1169	55.8	806	38.5	119	5.7
*Avoiding unnecessary travel during spread of Covid-19	1180	56.4	632	30.2	282	13.0
*Avoiding visiting relatives during spread of Covid-19	1523	72.7	408	19.5	163	7.8
* Using specialized personalized equipment	1288	61.5	806	38.5	0	0.0
*Using the appropriate methods in hand-washing	1077	51.4	982	46.9	35	1.7
* Using the appropriate methods in mask-wearing	1494	71.3	600	28.7	0	0.0

Table 5: this table shows that 85.4% of studied students are always not found in crowded places, 11% occasional while 3.6% are rarely found. 59.2% of them always wear

facemasks outside home, 39.2% occasional and 1.6% never wear mask. Furthermore, 57.2% always clean their hands by antiseptic, 33.9 were occasional and only 5.7% never. Also, 55.8% of the studied students always use tissue papers to cover mouth and nose during sneezing or coughing, 38.5% occasional and 5.7% never. 72.7% always don't visit their relatives during spread of Covid-19, 19.5% occasional and 7.8% ever. In addition, 38.5% of the studied students occasionally use specialized personalized equipment, 46.9% of them occasionally use the appropriate methods in hand washing and 28.7% use the appropriate methods in mask-wearing.

Figure 2: Distribution of studied university students according to their total satisfactory & unsatisfactory preventive Measures (n=2094).




Figure 2: as regards total satisfactory preventive measures of university students this figure illustrated that 83.7% of the studied students had always, 15.3% of them had occasional and only 1.0 % of them never regarding adherence to the preventive measures regarding Covid-19 disease outside the home. While, unsatisfactory practices the results clarified that, 48.5% had always, 12.3% had occasional, and 39.2 never adherence to the preventive measures regarding Covid-19 disease.

Table 6: Distribution of university students according to their perception regarding Covid-19 disease (n=2094)

Items	Agree		Not agree		Don't know	
	N	%	N	%	N	%
* Agree that the governorate successfully controlled in elimination of Covid?	1287	61.5	586	28.0	220	10.5
*Social communication platforms are our fears	1188	56.7	735	35.1	171	8.2
* changed for someone if they are affected with Covid-19	1261	60.2	675	32.2	158	7.5
* Ask for consult from responsible authorities in the event of emergency symptoms	1694	80.9	62	3.0	338	16.1
* Think the local authorities will succeed in reducing the spread of disease	1237	59.1	718	34.3	139	6.6
* Think the preventive measures of the government were early-produced	1359	64.9	652	31.1	83	4.0
* Think the preventive measures of the government were enough	1424	68.0	549	26.2	121	5.8
* Think the Covid-19 will last for a long time	1126	53.8	826	39.4	142	6.8
* Not necessary to take precautionary measures for children and older	1240	59.2	682	32.6	172	8.2
*Isolation of affected people is an effective way to reduce the spread of the Covid-19	1240	59.2	590	28.2	264	12.6
* Isolation and early treatment are the effective ways to control the prevalence of Covid-19	1221	58.3	676	32.3	197	9.4

Table 6: demonstrates that 61.5% of the studied students agreed that Palestinian governorate successfully controlled the in-elimination of covid-19, 28% do not agreed and 10.5% do not know, 56.7% agreed that social communication platforms are our fears, a higher percentage of students 80.9% would seek assistance from the authorities' responsible if the person or a family member develops symptoms of covid-19. 68% of students considered that the preventative measures taken by the Palestinian government at the beginning were sufficient and on a timely manner. 53.8% were worried toward covid-19. 59.2% of them remarkably agree that precautionary measures for children and older are not necessary & thought that covid-19 preventative measures should be applied by everyone to reduce spread the covid-19. In addition, 58.3% of the students would seek assistance from the authorities' responsible if the person or a family member develops symptoms of covid-19.

Table 7: Correlation between total knowledge, preventive measures and perception regarding Covid-19 disease (n=2094)

Items	Total students' knowledge				χ ²	P
	Satisfactory (n=1890)		Unsatisfactory (n=204)			
	No.	%	No.	%		
Total preventive measures outside home						
Always 80.2% (n=1680)	1581	83.7	99	48.5	63.73	<0.001* *
Occasional 15% (n= 315)	290	15.3	25	12.3		
Never 4.8% (n=99)	19	1.0	80	39.2		
Total perception of studied students						
Positive 78.3% (n=1640)	1460	77.2	180	88.2		
Negative 21.7% (n= 454)	430	22.8	24	11.8		

* Statistically significant P < 0.05 ** highly statistically significant P < 0.001

Table 7: there was a statistically highly significant relation between total knowledge and total practices for preventive measure outside home P-value<0.001, and statistically significant relation between total knowledge and total perception at P-value 0.005.

DISCUSSION:

University students play an important role in the community. During the Covid-19 pandemic crisis, university students are expected to spread attentiveness of key health and hygiene messages amongst communities. Even staying at home, retaining social distance, wearing face masks, washing hands, etc. are quite a few measures that the governments' health department propagate. So, it is so important that university students across the country have many knowledge and perception about all aspects of the disease including prevention strategies [19].Therefore, the study aimed to assess the extent of adherence to the Precautionary measures among University Students in Palestine during COVID-19 Pandemic.

Regarding socio-demographic characteristics of the studied university students, the results of the current study showed that, near to one third of them were in age group 17 to less than 19 years; with the average age 19.56 ± 3.069years. This finding was similar with the result of [13].in the study about the impact of the Covid-19 epidemic on mental health of undergraduate students in New Jersey, who reported that average age was 19.18 ± 2.9 years old. The finding of the present study illustrated that more than half of studied students were male students, this finding was unsupported by [19].found that more than three quarter of the studied students were female.

In accordance to knowledge of university students about Covid-19, the current study showed that the great majority of the studied students have correct answers about the meaning of covid-19, its causes & the treatment. All of them have correct answers about transmission of Covid; also, the majority of them have correct answers regarding signs and symptoms & protection methods from infection. A similar study conducted on a

Chinese population reported an overall knowledge of 90% [27]. The same table also reported 56.3% of students only know the time for repetition tests, more than one-tenth has incomplete answers, and more than one-third did not know the time. Regarding test to discover the virus & anticoagulation drugs showed that more than three-third of university student have correct answers. In addition, more than two-third have correct answers about timing to give anticoagulation. This result agrees with another study conducted on an Egyptian population comprising 559 participants reported the mean and standard deviation of knowledge score as 16.39 ± 2.63 , ranging from 7 to 22, which corresponds to approximately 74.5% overall knowledge among participants regarding Covid-19 [3]. Concerning to knowledge of university students according isolation the patients with Covid-19 at home this result reveals that the great majority of studied students have correct answers regarding the determination of the patient who needs isolation at home, meaning of home isolation, types of medication used during isolation, precaution for visitors, and precautions measures for patients with Covid-19 isolation. Furthermore, environmental cleaning & with 80.2% of the studied students had a satisfactory total knowledge regarding followed the precaution measures outside home. Our findings may be the current state of public health awareness and to determine the need for proper dissemination of knowledge and awareness. In addition, the easy access to highly recommended mass media, medical articles and journals available at the university's library which is the major source of information about Covid19 epidemics.

Regarding knowledge of university students followed preventive measures outside home, the current study found that the majority of students had a correct answer regarding following the preventive measures outside home and negligible number of them had incorrect answers regarding followed the preventive measures for respiratory hygiene to protect from Covid-19. Also, great number of them has correct answers regarding using personal protective equipment. This result was confirmed with [11], who studied knowledge, perception and preventive measures practices towards Covid-19, and found positive and cautious knowledge towards the Covid-19 epidemic were the vast majority of participants knew the precaution measures for respiratory hygiene and using the personal protective equipment, and avoid contact with others during the rapid rise period of the Covid-19.

In accordance to students' practices for preventive measures regarding Covid-19 disease outside home, the current study denoted that majority of them always are not found in crowded places, two-third always wear the face mask outside home, Furthermore, more than half of them always clean their hands with antiseptic solutions. Also, more than half of the studied students always use tissue papers to cover mouth and nose during sneezing or coughing, near to three quarter of them always do not visit their relatives during spread of Covid-19. In addition, more than one third of the studied students occasionally use specialized personalized equipment. This result agrees with [14]. who said in his study, throughout Covid-19 outbreak, results have shown that two-third of participants did not wear face masks when out in public, whereas fifth did not maintain physical distancing. Compliance with the pandemic restrictions is essential and crucial in

this phase, and failure to adhere to preventive measures-even if by a minority-would only lead to uncontrolled spreading of the disease. Positive practices towards Covid -19 pandemic were found to be significantly associated with the gender and major of the precipitant. This suggests may be that the students have a slightly positive preventive measures and perception regarding Covid -19.

Concerning the perception of university students among Covid-19 pandemic disease, the current study showed that more than two third of the sample agreed that Palestinian governorate successfully controlled in-elimination of covid-19, more than quarter did not agree, approximately two third agreed that social communication platforms are our fears, a higher percentage of students would seek assistance from the authorities' responsible if the person or a family member develops symptoms of Covid-19. 68% of students considered that the preventative measures taken by the Palestinian government at the beginning were sufficient and on a timely manner. More than half were worried toward \ covid-19 and two third of them remarkably thought that precautionary measures for children and older are not necessary & thought that covid-19 preventative measures should be applied by everyone to reduce spread the covid-19. As for the preventive measures applied by the government, showed that approximately two third of the students would seek assistance from the authorities' responsible if the person or a family member develops symptoms of covid-19. Concerning the performance of the local authorities, two thirds of participants considered the authorities' attempts successful in controlling the spread of Covid-19, and around half of them thought that the preventative measures were sufficient. Furthermore, less than a quarter of the participants believed that local authorities have sufficient tools to deal with the suspected cases of Covid-19. These findings disagree with other studies among different communities [27]; [18]; [4], that might be due to the lack of economic and medical resources and the shortage within the health system compared to developed countries. This indicates the importance of continuous health education that could improve the prevention behaviour toward Covid-19 in society.

CONCLUSION:

Based on the current study, the university students have substantial knowledge, practices for preventive measures, and a positive perception toward Covid -19. A majority of them also expressed their optimism regarding the control of Covid -19. In addition, there was a statistically highly significant relation between total knowledge and total practices outside home p -value <0.001 , and statistically significant relation between total knowledge and total perception p -value 0.005.

Recommendation:

- Continuing Government programs should aim to educate individuals from other sectors of the society to ensure the proper dissemination of knowledge on preventive safety measures, as this will help restrict and control the pandemic.
- The primary action to create awareness among the students can be undertaken through social media, electronic media and other possible means.

- It is the responsibility of the state to take all preventive measures for the safety of the community and public. There is an urgent need for research to address burden of the Covid-19 pandemic on college students.

Competing interests:

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Authors' contributions:

Each author took part in the design of the study, contributed to data collections, participated in writing the manuscript and the authors agree to accept equal responsibility for the accuracy of this paper. All authors approved the final article.

Acknowledgments:

We would like to thank all university students in Palestine who agreed to participate in the study.

References:

1. **Abdelhafiz AS, Mohammed Z, Ibrahim ME, Ziady HH, Alorabi M, Ayyad M (2020):** Knowledge, Perceptions, and Attitude of Egyptians Towards the Novel Coronavirus Disease (COVID-19). *J Community Health*. 2020; 1-10. doi: 10.1007/s10900-020-00827-7.
2. **Ahmed, A. (2020):** Knowledge, attitude and practices (KAP) towards COVID-19 and assessment of risks of infection by SARS-CoV-2 among the Bangladeshi population: An online cross sectional survey. <http://dx.doi.org/10.21203/rs.3.rs-24562/v2>.
3. **Azlan, A. Hamzah, MR, and Sern, TJ, (2020):** Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PLOS ONE* 15:e0233668. doi: 10.1371/journal.pone.0233668. doi: 10.1371/journal.pone.0233668. eCollection 2020.
4. **Bialek, S. Boundy, E. Bowen, V. (2020):** Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) — United States, February 12 March 16, 2020. *MMWR Morbidity and Mortality Weekly Report* 69:343–346. doi: 10.15585/mmwr.mm6912e2. DOI: <http://dx.doi.org/10.15585/mmwr.mm6912e2external icon>
5. **Chen, N. Zhou, M. and Dong, X. (2020):** Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *www.thelancet.com* 395:507. doi:10.1016/S0140-6736(20)30211-7.
6. **Elhadi, M. Msherghi, A. Alsoufi, A. Buzreg, A. Bouhuwaish, A. Khaled, A. Alhadi, A. Alameen, H. Biala, M. Elgherwi, E. Elkhafeefi, F. Elmabrouk, A. Abdulmalik, A. Alhaddad, S. Khaled, A. Elgzairi, M. (2020):** Knowledge, preventive behavior and risk perception regarding COVID-19: a self-reported study on college students Pan Africa Medical Journal. 2020;35(2):75 (ISSN: 1937-8688) <https://www.researchgate.net/publication/342535052>. DOI: 10.11604/pamj.supp.2020.35.2.23586
7. **Escalera, JP. Lizon, NF. Maldonado, A, (2020):** Clinical features of cases and a cluster of Coronavirus Disease 2019 (COVID-19) in Bolivia imported from Italy and Spain. *Travel Medicine and Infectious Disease* 101653. doi: 10.1016/j.tmaid.2020.101653.
8. **Hani, A. Naseef, N. Al-Shami, S. Abu Hadba, A. Leen, A. Humos, A. (2020):** Knowledge, attitudes and practices about Coronavirus disease (COVID-19) among Birzeit University students: A cross-sectional study, DOI: <https://doi.org/10.21203/rs.3.rs-83268/v1>, research square.
9. **Haque T, Hossain KM, Bhuiyan MMR, Ananna SA, Chowdhury SH, (2020):** Knowledge, attitudes, and practices towards COVID-19 among the Palestinian community of immense importance and is

critical for the community prevention/control of this Pandemic. <https://doi.org/10.1186%2Fs12889-021-10575-5>

10. **Hussain A, Garima T, Singh B, Ram R, Tripti R. (2020):** Knowledge, attitudes, and practices towards COVID-19 among Nepalese Residents: A quick online cross-sectional survey. *Asian J Med Sci.* January 5 2020;11: 6-11. <https://doi.org/10.3126/ajms.v11i3.28485>
11. **Novel Coronavirus (2019-nCoV) Situation report-72, 1st April 2020.** Geneva, Switzerland, 2020. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports> [Accessed 10th March 2020].
12. **Keckojevic, A. Basch, CH., Sullivan, M. Davi, NK. (2020):** The impact of the COVID-19 epidemic on mental health of undergraduate students in New Jersey, cross-sectional study. *PLoS ONE* 15(9): e0239696. <https://doi.org/10.1371/journal.pone.0239696>
13. **Khader, Y. Al Nsour, M. Al-Batayneh, OB. (2020):** Dentists' Awareness, Perception, and Attitude Regarding COVID-19 and Infection Control: Cross-Sectional Study Among Jordanian Dentists. *JMIR Public Health and Surveillance* 6:e18798. doi: 10.2196/18798
14. **Kumar J., Katto M. S., Siddiqui A. A., Sahito B., Jamil M., Rasheed N. and Ali M. (2020).** Knowledge, Attitude, and Practices of Healthcare Workers Regarding the Use of Face Mask to Limit the Spread of the New Coronavirus Disease (COVID-19). *Cureus.* 12 (4): 1-8. doi: 10.7759/cureus.7737.
15. **Olum R., Chekwech G., Wekha G., Nassozi D. R. and Bongomin F. (2020).** Coronavirus Disease-2019: Knowledge, Attitude, and Practices of Health Care Workers at Makerere University Teaching Hospitals, Uganda. *Frontiers in Public Health.* 8:181. DOI: 10.3389/fpubh.2020.00181.
16. **Palumbo R. (2017).** Examining the impacts of health literacy on healthcare costs. An evidence synthesis. *Health services management research.* 30(4):197-212.
17. **Rugarabamu, S. Byanaku, A. Ibrahim, M. (2020):** Knowledge, attitudes, and practices (KAP) towards COVID-19: A quick online cross-sectional survey among Tanzanian residents. *medRxiv* 2020.04.26.20080820. doi: 10.1101/2020.04.26.200808
18. **Reddy, P. & Srivastava, B. (2020):** Assessment of KAP (knowledge, attitude and practice) of university students towards prevention of Covid-19. *International Journal of Biological Innovations IJBI* 2 (2), (DECEMBER 2020) 117-125 <http://ijbi.org.in> | <http://www.gesa.org.in/journals.php>
19. **Virlée J., van Riel A. C. and Hammedi W. (2020):** Health literacy and its effects on wellbeing: how vulnerable healthcare service users integrate online resources. *Journal of Services Marketing.* (1) (34): 1-10.
20. **Wang, D. Hu, B. Hu, C. Zhu, F. Liu, X. Zhang, J. (2020):** Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA.* 2020; 323(11):1061–9. [PubMed] DOI: 10.1001/jama.2020.1585.
21. **White, MS. Omer, DM. Mohammad, DG. (2020):** Knowledge, Attitude and Practice on Prevention of Airborne and Droplet Infections during the Outbreak of Corona Virus among the College Students in University of Bisha, Saudi Arabia. *International Journal of Contemporary Research and Review* 11:20773–20776–20773–20776. doi: 10.15520/IJCRR.V11I04.802
22. **World Health Organization Coronavirus disease 2019 (COVID-19):** Situation Report – 91. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200420-sitrep-91-covid-19.pdf?sfvrsn=fcf0670b_4. Accessed 8 May 2020

23. **WHO Novel Coronavirus (2019- nCoV) Situation report-19, 8 February (2020)** Geneva, Switzerland, 2020. Available from: <https://www.who.int/emergencies/diseases/novelcoronavirus-2019/situation-reports> [Accessed 10th March 2020].
24. **WHO /Europe (2020):** Coronavirus disease (COVID-19) outbreak - WHO announces COVID-19 outbreak a pandemic. Announces -covid-19-outbreak-a-pandemic. Accessed 8 Sep 2020.<https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who>
25. **Zhang, M. Zhou, M. Tang, F. (2020):** Knowledge, attitude, and practice regarding COVID-19 among healthcare workers in Henan, China. *Journal of Hospital Infection* 105:183–187. doi: 10.1016/j.jhin.2020.04.012.
26. **Zhong BL, Luo W, Li HM, Zhang QQ, Liu XG, Li WT (2020):** Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci.* 2020; 16(10):1745-1752. 19. doi: 10.7150/ijbs.45221.
27. **ELBASHA, M. 2020.** Palestinians confirm 7 coronavirus cases, declare tourist ban [Online]. Available: <https://www.timesofisrael.com/palestinians-confirm-7-coronavirus-cases-declare-tourist-ban/> [Accessed].
28. **ALSER, O., ALWAHEIDI, S., ELESSI, K. & MEGHARI, H. 2020. COVID-19 in Gaza:** a pandemic spreading in a place already under protracted lockdown. *Eastern Mediterranean Health Journal*, 26, 762-763.
29. **UNRWA. 2019.** WHERE WE WORK [Online]. Available: <https://www.unrwa.org/where-we-work/gaza-strip> [Accessed].
30. **The Coalition for Epidemic Preparedness Innovations (CEPI) (2020).** CEPI welcomes UK Government's funding and highlights need for \$2 billion to develop a vaccine against COVID-19. 2020. Available from: https://cepi.net/news_cepi/2-billion-required-to-develop-a-vaccine-against-the-covid-19-virus. [cited 2020 Mar 23].
31. **.Michie S, West R. Behavioural, (2020):** environmental, social, and systems interventions against covid-19. *BMJ.* 2020;370. Available from: <https://www.bmj.com/content/370/bmj.m2982>.
32. **Kang L et al.,(2020).** Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: a cross-sectional study. *Brain Behav Immun* 87: 11–17. <https://doi.org/10.1016%2Fj.psychres.2020.113190>